## \*\*11/12/04 DRAFT\*\*

# Fire Regime Condition Class (FRCC) Interagency Handbook Reference Conditions

Modeler: Wendel Hann, Reese Lolley, Date: 10/08/04 PNVG Code: PMGR2

Cecilia McNicoll

Potential Natural Vegetation Group: Plains Mesa Grassland with Trees

**Geographic Area:** Southeastern Arizona and southwestern New Mexico.

**Description**: The Plains Mesa Grassland is described by Brown (1994) as occurring on open and exposed plains and the intermingled foothills subject to high heat and wind. The Plains Mesa Grassland with Tree (PMGR2) occurs at mid- to upper elevations (5000-7500 ft) in southern Arizona and New Mexico, in a zone above or in a mosaic with Madrean Evergreen Woodland (MAME), sometimes chaparral (CHAP5), and often adjacent to the Ponderosa Pine (PPIN) zone at upper elevations. In its southern range, PMGR2 occurs in a zone above desert grassland (DGRA2). PMGR2 have mild winters and hot, windy summers with precipitation ranging from about 12 to 18 inches. Almost half the precipitation occurs in the summer, with the other half occurring in winter and spring. In the northern extension and at higher elevations, were there is a stronger Great Basin influence, cooler summer and winter temperatures occur with overall less precipitation received predominantly in the winter and spring. Landforms typically include mesas, foothills, and mountains. Kuchler (1964) did not identify this type, but the closest fit to the east would be the Grama-Buffalo Grass (Type 65), and Gramma-tobosa Shrubsteppe (58) to the north. (In a previous version of the FRCC coarse-scale models, this vegetation type was included in the Plains Grassland [PGRA] model).

\*Note: This PNVG is somewhat broader than the Plains Mesa Grassland type defined by Dick-Peddie (1993)(i.e., includes a portion of that classification's Woodland and Savanna type).

**Fire Regime Description**: Fire Regime II (frequent replacement). Presettlement fire intervals likely ranged from 3-10 years long, with an estimated MFI of 5 years (Brown and Smith 2000, USDA 2002). Frequent fires promoted grass dominance, and inhibited associated shrubs and trees. Fires generally caused total replacement of the upper vegetation layer, followed by resprouting grasses with the onset of seasonal precipitation.

### **Vegetation Type and Structure**

Class	Percent of		
	Landscape		
A: Early -seral	2	Post-fire grass and fire-adapted forbs	
B: Mid-seral Closed	30	Mid-seral productive closed grass on better soils and	
Grassland		shifting from class C to B in moist years.	
C: Mid-seral Open	60	Mid-seral less productive open grass on lower moisture	
Grassland		holding soils and shifting from class B to C in drought years.	
D: Late-seral Open	7	Late-seral open tree and shrub communities with grass	
Tree-Shrub/Grass		understory and patches between trees where topo- edaphic and micro site conditions reduce fire frequency	

E: Late-seral Closed Tree/Shrub

or severity so that trees and shrubs survive.

Late-seral closed tree and shrub communities topoedaphic and micro site conditions reduce fire frequency

or severity so that trees and shrubs can survive.

Total 100

Fire Frequency and Severity

ine inequency and bevenly					
Fire Frequency-	Modeled	Percent,	Description		
Severity	Probability	All Fires			
Replacement Fire	.19	99	Replacement fires in A, B and C		
Non-Replacement Fire	.01	1	Mosaic fires in classes D and E		
All Fire Frequency*	.20	100	5-yr MFI overall (generally low variation)		

<sup>\*</sup>Sum of replacement fire and non-replacement fire probabilities.

#### References

Brown, David E., editor. 1982. Biotic communities of the American Southwest–United States and Mexico. Desert Plants 4(1-4): 1-342.

Brown, James K.; Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Dick-Peddie, W. A., 1993. New Mexico vegetation, past, present, and future. University of New Mexico Press.

Kuchler, A. W. 1964. Manual to accompany the map of potential natural vegetation of the conterminous United States. American Geographical Society. Spec. Publ. No. 36. Lib. Congress Cat. Card Num. 64-15417. 156 p.

Schmidt, Kirsten M, Menakis, James P., Hardy, Colin C., Hann, Wendel J., Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.

U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online]. Available: <a href="http://www.fs.fed.us/database/feis/">http://www.fs.fed.us/database/feis/</a>.

#### MODELER FIELD REVIEWS:

Wendel Hann - Colorado 2003, New Mexico 2003, Texas 2001

## **VDDT Results:**











